


**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

30

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 401/14, A61K 31/395</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/47573</b> <b>(43) International Publication Date:</b> 17 August 2000 (17.08.00)
<b>(21) International Application Number:</b> PCT/GB00/00354 <b>(22) International Filing Date:</b> 8 February 2000 (08.02.00) <b>(30) Priority Data:</b> 9902989.4 11 February 1999 (11.02.99) GB <b>(71) Applicant (for all designated States except US):</b> AS-TRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). <b>(72) Inventor; and</b> <b>(75) Inventor/Applicant (for US only):</b> CAULKETT, Peter, William, Rodney [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). <b>(74) Agent:</b> BROWN, Andrew, Stephen; Global Intellectual Property, AstraZeneca UK Limited, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).	<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> With international search report.	

**(54) Title:** HETEROCYCLIC DERIVATIVES AS INHIBITORS OF FACTOR XA**(57) Abstract**

The invention relates to heterocyclic derivatives, or pharmaceutically-acceptable salts thereof, of formula (I) wherein: A is an optionally substituted 5- or 6-membered monocyclic aromatic ring containing 1, 2 or 3 ring heteroatoms selected from oxygen, nitrogen and sulphur atoms; which possess antithrombotic and anticoagulant properties and are accordingly useful in methods of treatment of humans or animals. The invention also relates to processes for the preparation of the heterocyclic derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments for use in the production of an antithrombotic or anticoagulant effect.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## HETEROCYCLIC DERIVATIVES AS INHIBITORS OF FACTOR XA

The invention relates to heterocyclic derivatives, or pharmaceutically-acceptable salts thereof, which possess antithrombotic and anticoagulant properties and are accordingly  
5 useful in methods of treatment of humans or animals. The invention also relates to processes for the preparation of the heterocyclic derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments for use in the production of an antithrombotic or anticoagulant effect.

The antithrombotic and anticoagulant effect produced by the compounds of the  
10 invention is believed to be attributable to their strong inhibitory effect against the activated coagulation protease known as Factor Xa. Factor Xa is one of a cascade of proteases involved in the complex process of blood coagulation. The protease known as thrombin is the final protease in the cascade and Factor Xa is the preceding protease which cleaves prothrombin to generate thrombin.

15 Certain compounds are known to possess Factor Xa inhibitory properties and the field has been reviewed by R.B. Wallis, Current Opinion in Therapeutic Patents, 1993, 1173-1179. Thus it is known that two proteins, one known as antistatin and the other known as tick anticoagulant protein (TAP), are specific Factor Xa inhibitors which possess antithrombotic properties in various animal models of thrombotic disease.

20 It is also known that certain non-peptidic compounds possess Factor Xa inhibitory properties. Of the low molecular weight inhibitors mentioned in the review by R.B. Wallis, all possessed a strongly basic group such as an amidinophenyl or amidinonaphthyl group.

We have now found that certain heterocyclic derivatives possess Factor Xa inhibitory activity. Many of the compounds of the present invention also possess the  
25 advantage of being selective Factor Xa inhibitors, that is the enzyme Factor Xa is inhibited strongly at concentrations of test compound which do not inhibit or which inhibit to a lesser extent the enzyme thrombin which is also a member of the blood coagulation enzymatic cascade.

The compounds of the present invention possess activity in the treatment or  
30 prevention of a variety of medical disorders where anticoagulant therapy is indicated, for example in the treatment or prevention of thrombotic conditions such as coronary artery and

- 2 -


cerebro-vascular disease. Further examples of such medical disorders include various cardiovascular and cerebrovascular conditions such as myocardial infarction, the formation of atherosclerotic plaques, venous or arterial thrombosis, coagulation syndromes, vascular injury including reocclusion and restenosis following angioplasty and coronary artery bypass

5 surgery, thrombus formation after the application of blood vessel operative techniques or after general surgery such as hip replacement surgery, the introduction of artificial heart valves or on the recirculation of blood, cerebral infarction, cerebral thrombosis, stroke, cerebral embolism, pulmonary embolism, ischaemia and angina (including unstable angina).

The compounds of the invention are also useful as inhibitors of blood coagulation in  
10 an ex-vivo situation such as, for example, the storage of whole blood or other biological samples suspected to contain Factor Xa and in which coagulation is detrimental.

Accordingly in one aspect the present invention provides compounds of the formula

I



15

wherein:

A is an optionally substituted 5- or 6-membered monocyclic aromatic ring containing 1, 2 or 3 ring heteroatoms selected from oxygen, nitrogen and sulphur atoms;

B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> are independently CH or a nitrogen atom, wherein the ring formed from B<sub>1</sub>,

20 B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> may optionally be substituted; with the proviso that at least one of B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is nitrogen;

T<sub>1</sub> is CH or N;

T<sub>2</sub> is CH or N; with the proviso that at least one of T<sub>1</sub> and T<sub>2</sub> is N;

X<sub>1</sub> is SO, SO<sub>2</sub>, C(R<sub>4</sub>)<sub>2</sub> or CO when T<sub>1</sub> is CH or N; or in addition X<sub>1</sub> is O or S when T<sub>1</sub> is CH;

25 and wherein each R<sub>4</sub> is independently hydrogen or (1-4C)alkyl;

L<sub>1</sub> is (1-4C)alkylene or (1-3C)alkylenecarbonyl;

R<sub>2</sub> is hydrogen or (1-4C)alkyl;

R<sub>3</sub> is hydrogen or (1-4C)alkyl;

- 3 -

or  $R_2$  and  $R_3$  are joined to form a (1-4C)alkylene or  $-\text{CH}_2\text{CO}-$  group; wherein the ring formed by  $T_1$ ,  $R_2$ ,  $R_3$ ,  $T_2$  and  $L_1$  is optionally substituted;

$X_2$  is  $\text{S(O)}_y$  wherein  $y$  is one or two,  $\text{C(R}^5)_2$  or  $\text{CO}$ ; and each  $\text{R}^5$  is independently hydrogen or (1-4C)alkyl;

- 5 Y is selected from hydrogen, halo, trifluoromethyl, trifluoromethoxy, cyano, hydroxy, amino, nitro, carboxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-(1-4C)alkylamino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl, (2-4C)alkanoyl, (2-4C)alkanoylamino, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, carboxy-(1-4C)alkyl, (1-4C)alkoxycarbonyl-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-(1-4C)alkyl and N,N-di-(1-4C)alkylcarbamoyl-(1-4C)alkyl;
- 10  $n$  is 1 or 2; and

$B_5$  and  $B_6$  is selected from N or CH; with the proviso that at least one of  $B_5$  and  $B_6$  is N;

- 15 and pharmaceutically acceptable salts thereof.

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. An analogous convention applies to other generic terms.

- It is to be understood that certain heterocyclic derivatives of the present invention can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms which possess Factor Xa inhibitory activity.
- 20

- It is further to be understood that, insofar as certain of the compounds of the formula defined above may exist in optically active or racemic forms by virtue of one or more asymmetric carbon atoms, the invention encompasses any such optically active or racemic form which possesses Factor Xa inhibitory activity. The synthesis of optically active forms may be carried out by standard techniques of organic chemistry well known in the art, for example by synthesis from optically active starting materials or by resolution of a racemic form.
- 25

- 30 Preferably A is a pyridyl, pyrimidinyl or pyridazinyl ring for example 4-pyridyl,

- 4 -

2-pyridyl, 4-pyridazinyl, 3-pyrimidinyl, 4-pyrimidinyl or 3-pyridyl. Of these 4-pyrimidinyl, 4-pyridazinyl and 4-pyridyl are preferred, with 4-pyrimidinyl and 4-pyridyl most preferred.

In one aspect A is unsubstituted. In another aspect A is substituted by one, two or three atoms or groups selected from halo (for example fluoro, chloro or bromo),  
 5 trifluoromethyl, cyano, amino, oxo, hydroxy, nitro, (1-4C)alkyl (for example methyl or ethyl), (1-4C)alkoxy (for example methoxy or ethoxy), (1-4C)alkylamino (for example methylamino or ethylamino) or di-(1-4C)alkylamino (for example dimethylamino or diethylamino). For the avoidance of doubt substituents may also be on any heteroatom.

Preferably the ring formed by B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is a pyridinediyl, wherein B<sub>1</sub>, or B<sub>3</sub>  
 10 is a nitrogen atom, pyrimidinediyl, wherein B<sub>1</sub> and B<sub>2</sub> or B<sub>3</sub> and B<sub>4</sub> are nitrogen atoms, pyridazinediyl, wherein B<sub>1</sub>, B<sub>3</sub> and B<sub>4</sub> or B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub> are nitrogen atoms. Of these pyridinediyl and pyrimidinediyl are preferred, and pyridinediyl is most preferred.

In one aspect the ring containing B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is unsubstituted. In another aspect the ring containing B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is substituted by one or two  
 15 substituents selected from hydroxy, carboxy, (1-4C)alkoxycarbonyl or one of the following;

$-(CH_2)_n-R$ ,  $-(CH_2)_n-NRR_1$ ,  $-CO-R$ ,  $-CO-NRR_1$ ,  $-(CH_2)_n-CO-R$  and  $-(CH_2)_n-CO-NRR_1$ ;

wherein n is 1 or 2;

20 R and R<sub>1</sub> are independently selected from hydrogen, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, hydroxy(1-4C)alkyl, carboxy(1-4C)alkyl and (1-4C)alkoxycarbonyl-(1-4C)alkyl or where possible R and R<sub>1</sub> may together form a 5- or 6-membered optionally substituted heterocyclic ring which may include in addition to the nitrogen atom to which R and R<sub>1</sub> are attached 1 or 2 additional heteroatoms selected from nitrogen, oxygen and sulphur.

25 In a particular aspect the heterocyclic rings formed by R and R<sub>1</sub> are preferably selected from pyrrolidin-1-yl, imidazolin-1-yl, piperidin-1-yl, piperazin-1-yl, 4-morpholino and 4-thiomorpholino. In a particular aspect the heterocyclic ring formed by R and R<sub>1</sub> may be unsubstituted. In an alternative aspect the ring formed by R and R<sub>1</sub> is substituted by 1 or 2 substituents selected from oxo, hydroxy and carboxy.

30 In a particular aspect, when T<sub>1</sub> is CH or N, X<sub>1</sub> is CO, SO<sub>2</sub>, or CH<sub>2</sub> or, when T<sub>1</sub> is CH, X<sub>1</sub> in addition is O or S. Preferably X<sub>1</sub> is CO.

- 5 -

$T_1$  is CH or N and  $T_2$  is CH or N with the proviso that at least one of  $T_1$  and  $T_2$  is N. For the avoidance of doubt  $T_1$  is directly attached to the groups  $X_1$  and  $L_1$  and  $T_2$  is directly attached to the groups  $L_1$  and  $X_2$ .

$L_1$  is (1-4C)alkylene for example methylene, ethylene or propylene or is

5 C<sub>1-3</sub>alkylenecarbonyl for example methylenecarbonyl (-CH<sub>2</sub>CO-), preferably  $L_1$  is ethylene.

In one aspect  $R_2$  is hydrogen or (1-4C)alkyl for example methyl or ethyl. In one aspect  $R_3$  is hydrogen or C<sub>1-4</sub>alkyl for example methyl or ethyl.

In a preferred aspect  $R_2$  and  $R_3$  are joined to form a (1-4C)alkylene group, for example a methylene, ethylene or propylene group, or a methylenecarbonyl (-CH<sub>2</sub>CO-) group,  
10 preferably ethylene.

In a particular aspect  $R_2$  and  $R_3$  are joined to form, together with  $T_1$ ,  $T_2$  and  $L_1$ , a heterocyclic ring wherein at least one of  $T_1$  and  $T_2$  is N. Examples of such heterocyclic rings are piperazine (wherein  $T_1$  and  $T_2$  are both N), piperidine (wherein either  $T_1$  or  $T_2$  is N and the other is CH) and pyrrolidine (wherein either  $T_1$  or  $T_2$  is N and other is CH).

15 In one aspect the heterocyclic ring formed by  $T_1$ ,  $T_2$ ,  $L_1$ ,  $R_2$  and  $R_3$  is unsubstituted. In another aspect this ring is substituted by one or two substituents selected from hydroxy, oxo, carboxy, (1-4C)alkoxycarbonyl or one of the following;

20  $-(CH_2)_n-R$ ,  $-(CH_2)_n-NRR_1$ ,  $-CO-R$ ,  $-CO-NRR_1$ ,  $-(CH_2)_n-CO-R$  and  $-(CH_2)_n-CO-NRR_1$ ;

wherein  $n$  is 1 or 2;

$R$  and  $R_1$  are independently selected from hydrogen, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, hydroxy(1-4C)alkyl, carboxy(1-4C)alkyl and (1-4C)alkoxycarbonyl- (1-4C)alkyl or where possible  $R$  and  $R_1$  may together form a 5- or 6-membered optionally  
25 substituted heterocyclic ring which may include in addition to the nitrogen atom to which  $R$  and  $R_1$  are attached 1 or 2 additional heteroatoms selected from nitrogen, oxygen and sulphur.

In a particular aspect the heterocyclic rings formed by  $R$  and  $R_1$  are preferably selected from pyrrolidin-1-yl, imidazolin-1-yl, piperidin-1-yl, piperazin-1-yl, 4-morpholino and 4-thiomorpholino. In a particular aspect the heterocyclic ring formed by  $R$  and  $R_1$  may be  
30 unsubstituted. In an alternative aspect the ring formed by  $R$  and  $R_1$  is substituted by 1 or 2

- 6 -

substituents selected from oxo, hydroxy, carboxy and (1-4C)alkyl, preferably oxo, hydroxy, and carboxy.

In a particular aspect  $X_2$  is  $SO_2$ ,  $CH_2$  or CO. Preferably  $X_2$  is  $SO_2$ .

In a preferred aspect Y is selected from from hydrogen, halo (bromo or chloro),  
 5 trifluoromethyl, trifluoromethoxy, cyano, hydroxy, amino, nitro, carboxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-(1-4C)alkylamino and (1-4C)alkoxycarbonyl.

Suitable values for substituents Y are:

- |    |  |  |
|----|--|--|
| 10 | for halo:                                  | fluoro, chloro, bromo;   |
|    | for (1-4C)alkyl:                           | methyl, ethyl, propyl, butyl;  |
|    | for (1-4C)alkoxy:                          | methoxy, ethoxy;   |
|    | for (1-4C)alkylamino:                      | methylamino, ethylamino;   |
| 15 | for di-(1-4C)alkylamino:                   | dimethylamino, diethylamino;   |
|    | for (2-4C)alkenyl:                         | vinyl and allyl;   |
|    | for (2-4C)alkynyl:                         | ethynyl and prop-2-ynyl;   |
|    | for (2-4C)alkenyloxy:                      | vinylloxy and allyloxy;  |
|    | for (2-4C)alkynyloxy:                      | ethynyloxy and prop-2-ynyloxy;   |
| 20 | for (1-4C)alkylthio:                       | methylthio, ethylthio and propylthio;  |
|    | for (1-4C)alkylsulphinyl:                  | methylsulphinyl, ethylsulphinyl and propylsulphinyl;                               |
|    | for (1-4C)alkylsulphonyl:                  | methylsulphonyl, ethylsulphonyl and propylsulphonyl;                               |
| 25 | for (2-4C)alkanoylamino:                   | acetamido, propionamido and butyramido;  |
|    | for (1-4C)alkoxycarbonyl:                  | methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl and <u>tert</u> -butoxycarbonyl;  |
|    | for <u>N</u> -(1-4C)alkylcarbamoyl:        | <u>N</u> -methylcarbamoyl, <u>N</u> -ethylcarbamoyl and <u>N</u> -propylcarbamoyl; |
| 30 | for <u>N,N</u> -di-[(1-4C)alkyl]carbamoyl: | <u>N,N</u> -dimethylcarbamoyl, <u>N</u> -ethyl- <u>N</u> -methylcarbamoyl and      |

- 7 -

- for (2-4C)alkanoyl:  
 for hydroxy-(1-4C)alkyl:  
 5 for (1-4C)alkoxy-(1-4C)alkyl:  
 for carboxy-(1-4C)alkyl:  
 10 for (1-4C)alkoxycarbonyl-(1-4C)alkyl:  
 15 for carbamoyl-(1-4C)alkyl:  
 20 for N-(1-4C)alkylcarbamoyl-(1-4C)alkyl:  
 25 for N,N-di-[(1-4C)alkyl]carbamoyl-(1-4C)alkyl:  
 30
- N,N-diethylcarbamoyl;  
 acetyl, propionyl and butyryl;  
 hydroxymethyl, 1-hydroxyethyl,  
 2-hydroxyethyl and 3-hydroxypropyl;  
 methoxymethyl, ethoxymethyl,  
 1-methoxymethyl, 2-methoxyethyl,  
 2-ethoxyethyl and 3-methoxypropyl;  
 carboxymethyl, 1-carboxyethyl,  
 2-carboxyethyl and 3-carboxypropyl;  
 methoxycarbonylmethyl,  
 ethoxycarbonylmethyl, tert-butoxy-  
 carbonylmethyl, 1-methoxycarbonylethyl,  
 1-ethoxycarbonylethyl,  
 2-methoxycarbonylethyl,  
 2-ethoxycarbonylethyl,  
 3-methoxycarbonylpropyl and  
 3-ethoxycarbonylpropyl;  
 carbamoylmethyl, 1-carbamoylethyl,  
 2-carbamoylethyl and 3-carbamoylpropyl;  
N-methylcarbamoylmethyl,  
N-ethylcarbamoylmethyl,  
N-propylcarbamoylmethyl,  
 1-(N-methylcarbamoyl)ethyl,  
 1-(N-ethylcarbamoyl)ethyl,  
 2-(N-methylcarbamoyl)ethyl,  
 2-(N-ethylcarbamoyl)ethyl and  
 3-(N-methylcarbamoyl)propyl;  
N,N-dimethylcarbamoylmethyl,  
N-ethyl-N-methylcarbamoylmethyl,  
N,N-diethylcarbamoylmethyl,  
 1-(N,N-dimethylcarbamoyl)ethyl,

- 8 -

1-(N,N-diethylcarbamoyl)ethyl,  
 2-(N,N-dimethylcarbamoyl)ethyl,  
 2-(N,N-diethylcarbamoyl)ethyl and  
 3-(N,N-dimethylcarbamoyl)propyl;

5

A preferred class of compounds of the present invention is that wherein:

A is 4-pyridyl, 4-pyrimidinyl or 4-pyridazinyl;

B<sub>1 to 4</sub> is forms a pyridinediyl, pyrimidinediyl or pyridazinediyl;

X<sub>1</sub> is CO, SO<sub>2</sub> or CH<sub>2</sub>, ideally CO;

10 T<sub>1</sub> and T<sub>2</sub> are both N;

L<sub>1</sub> is ethylene or propylene;

R<sub>2</sub> and R<sub>3</sub> are joined to form an ethylene or propylene or methylenecarbonyl group;

wherein the heterocyclic ring formed by T<sub>1</sub>, T<sub>2</sub>, L<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> is unsubstituted or is substituted;

X<sub>2</sub> is SO<sub>2</sub>;

15 B<sub>5</sub> or B<sub>6</sub> is N:

n is 1 at the 5 position;

Y is halo, preferably bromo or chloro;

and pharmaceutically-acceptable salts thereof.

A particular compound of the invention is:

20 1-(5-chloroindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine; and  
 1-(5-bromoindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine.

Compounds of formula I, or pharmaceutically-acceptable salt thereof, may be prepared by any process known to be applicable to the preparation of related compounds.


Such procedures are provided as a further feature of the invention and are illustrated by the

25 following representative processes in which, unless otherwise stated A, B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub>, B<sub>4</sub>, X<sub>1</sub>, T<sub>1</sub>, T<sub>2</sub>, L<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, X<sub>2</sub>, B<sub>5</sub>, B<sub>6</sub>, Y and n have any of the meanings defined hereinbefore wherein any functional group, for example amino, alkylamino, carboxy or hydroxy, is optionally protected by a protecting group which may be removed when necessary.

Necessary starting materials may be obtained by standard procedures of organic  
 30 chemistry.

According to another aspect, the present invention provides a process for preparing a compound of formula I or a pharmaceutically acceptable salt thereof, which comprises:

- (a) For the production of compounds of the formula (I) wherein  $T_1$  is N and  $X_1$  is CO, by the reaction, conveniently in the presence of a suitable base, of an amine with an acid




or with a reactive derivative of the acid.

- 10 A suitable reactive derivative of the acid is, for example, an acyl halide, an anhydride, an activated amide, an active ester, or the product of the reaction of the acid and a carbodiimide such as N,N'-dicyclohexylcarbodiimide or N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide.

- The reaction is conveniently carried out in the presence of a suitable base such as, 15 for example, an alkali or alkaline earth metal carbonate, alkoxide, hydroxide or hydride, for example sodium carbonate, potassium carbonate, sodium ethoxide, potassium butoxide, sodium hydroxide, potassium hydroxide, sodium hydride or potassium hydride, or a dialkylamino-lithium, for example lithium di-isopropylamide, or, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, 20 triethylamine, morpholine or diazabicyclo[5.4.0]undec-7-ene. The reaction is also preferably carried out in a suitable inert solvent or diluent, for example methylene chloride, chloroform, carbon tetrachloride, tetrahydrofuran, 1,2-dimethoxyethane, N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidin-2-one, dimethylsulphoxide or acetone, and at a temperature in the range, for example, -78° to 150°C, conveniently at or near ambient 25 temperature.

- (b) For the production of those compounds of formula I wherein  $T_1$  is CH and  $X_1$  is O by the reaction, conveniently in the presence of a suitable coupling agent;

- 10 -



A suitable value for the displaceable group Z is, for example, a halogeno or sulphonyloxy group, for example a fluoro, chloro, bromo, mesyloxy or 4-tolylsulphonyloxy group.


- 5 A suitable reagent for the coupling reaction when Z is a halogeno or sulphonyloxy group is, for example, a suitable base, for example, an alkali or alkaline earth metal carbonate, hydroxide or hydride, for example sodium carbonate, hydroxide or hydride, for example sodium carbonate, potassium carbonate, sodium hydroxide, potassium hydroxide, sodium hydride or potassium hydride. The alkylation reaction is preferably performed in a  
 10 suitable inert solvent or diluent, for example N,N-dimethylformamide, N,N-dimethylacetamide, dimethylsulphoxide, acetone, 1,2-dimethoxyethane or tetrahydrofuran, and at a temperature in the range, for example, -10° to 150°C, conveniently at or near ambient temperature.

- An analogous procedure may be employed for the preparation of those compounds  
 15 of the formula (I) wherein T<sub>1</sub> is CH and X<sub>1</sub> is a group of the formula S.

- A suitable reagent for the coupling reaction of the alcohol, wherein Z is a hydroxy group, where the hydroxy group is converted in situ to a displaceable group as defined above, is, for example, the reagent obtained when said alcohol is reacted with a di-(1-4C)alkyl azodicarboxylate in the presence of a triarylphosphine or  
 20 tri-(1-4C)alkylphosphine, for example with diethyl azodicarboxylate in the presence of triphenylphosphine or tributylphosphine. The reaction is preferably performed in a suitable inert solvent or diluent, for example acetone, 1,2-dimethoxyethane or tetrahydrofuran, and at a temperature in the range, for example, 10° to 80°C, conveniently at or near ambient temperature.  
 25

- (c) For the production of those compounds of formula (I) wherein T<sub>1</sub> is N and X<sub>1</sub> is CH(R<sub>4</sub>), the reductive amination of a keto compound below:


- 11 -



with an amine as defined in (a) above.

A suitable reducing agent is, for example, a hydride reducing agent, for example  
 5 an alkali metal aluminium hydride such as lithium aluminium hydride or, preferably, an  
 alkali metal borohydride such as sodium borohydride, sodium cyanoborohydride, sodium  
 triethylborohydride, sodium trimethoxyborohydride and sodium triacetoxyborohydride. The  
 reaction is conveniently performed in a suitable inert solvent or diluent, for example  
 tetrahydrofuran and diethyl ether for the more powerful reducing agents such as lithium  
 10 aluminium hydride, and, for example, methylene chloride or a protic solvent such as  
 methanol and ethanol for the less powerful reducing agents such as sodium  
 triacetoxyborohydride. The reaction is performed at a temperature in the range, for example,  
 10° to 80°C, conveniently at or near ambient temperature.

15 (d) By the reaction of:



wherein Z is a displaceable group such as halo, with an activated derivative of heterocyclic  
 ring A. Suitable activated derivatives include metalised derivatives, such as with zinc or tin,  
 20 and borane derivatives. The activated derivative of heterocyclic ring A is reacted with the  
 above compound to effect cross coupling where Z is a halo group, such as iodo, bromo or  
 chloro and triflate. Suitably the reaction is catalysed by use of a transition state metal catalyst,  
 such as palladium, e.g. tetrakis (triphenylphosphine) palladium (0).


Alternatively it is possible that ring A contains the displaceable group Z and the ring  
 25 containing B<sub>1</sub> to B<sub>4</sub> is activated, as described above.

The reaction is not suitable for compounds which contain halo substituents on A, B,  
 or L<sub>1</sub>.

- (e) By forming A ring on the above compound (d), wherein Z is a functional group capable of cyclisation. Suitable reagents and conditions are described in Bredereck H. Chem.Ber.; 96, 1505, (1963); Fuchigami, T., Bull. Chem. Soc. Jpn., 49, p3607, (1976); Huffman, K.R., J. Org. Chem., 28, p1812, (1963); Palusso, G., Gazz. Chim. Ital., 90, p1290, (1960) and Ainsworth C.J., Heterocycl. Chem., 3, p470, (1966). Processes suitable for synthesis of starting materials in such cyclisation reactions are described in Zhang M.Q. et.al; J.Heterocyclic. Chem.; 28, 673, (1991) and Kosugi, M. et al., Bull. Chem. Soc. Jpn., 60, 767-768 (1987).


10

- (f) For the production of compounds wherein  $T_2$  is N, by the reaction:



- 15 wherein Z is a displaceable group for example chloro, under conditions similar to those of process variant (a) above.

- (g) For the production of compounds wherein  $T_1$  is N and  $X_1$  is SO or  $\text{SO}_2$ , by the reaction:



20

wherein x is one or two and Z is a displaceable group; under appropriate conventional coupling conditions, similar to those of process variant (a) above.

- (h) By coupling the heteroaryl group to  $T_2$  with methods analogous to those described in process variants (a), (c) and (f) for preparing the  $\text{B}-\text{X}_1-\text{T}_1$ - moiety may be employed.

25

- 13 -

(i) For the production of compounds of the formula (I) wherein  $X_1$  is a group of the formula SO,  $SO_2$ , wherein the ring containing  $B_1$  to  $B_4$  bears a 1-oxothiomorpholino or 1,1-dioxothiomorpholino group or a substituent which contains a (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, 1-oxothiomorpholino or 1,1-dioxothiomorpholino group, wherein  $X_2$  is a group of the formula SO or  $SO_2$ , wherein Q bears a (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, phenylsulphinyl, phenylsulphonyl, heteroarylsulphinyl or heteroarylsulphonyl group, the oxidation of the corresponding compound of the formula I wherein  $X_1$ ,  $X_2$ , or both  $X_1$  and  $X_2$  is S.

A suitable oxidising agent is, for example, any agent known in the art for the oxidation of thio to sulphinyl and/or sulphonyl, for example, hydrogen peroxide, a peracid (such as 3-chloroperoxybenzoic or peroxyacetic acid), an alkali metal peroxysulphate (such as potassium peroxymonosulphate), chromium trioxide or gaseous oxygen in the presence of platinum. The oxidation is generally carried out under as mild conditions as possible and with the required stoichiometric amount of oxidising agent in order to reduce the risk of over oxidation and damage to other functional groups. In general the reaction is carried out in a suitable solvent or diluent such as methylene chloride, chloroform, acetone, tetrahydrofuran or *tert*-butyl methyl ether and at a temperature, for example, at or near ambient temperature, that is in the range 15 to 35°C. Suitable reagents and conditions are described in, for example, Page G. O.; Synth. Commun. 23, (1993) 6, 765-769. When a compound carrying a sulphinyl group is required a milder oxidising agent may also be used, for example sodium or potassium metaperiodate, conveniently in a polar solvent such as acetic acid or ethanol. It will be appreciated that when a compound of the formula I containing a sulphonyl group is required, it may be obtained by oxidation of the corresponding sulphinyl compound as well as of the corresponding thio compound. Those compounds of formula I which contain oxygen labile groups (such as A ring is pyridyl) are probably not suitable intermediates for this process step, unless oxidation of such groups is desired.

When a pharmaceutically-acceptable salt of a compound of the formula I is required, it may be obtained, for example, by reaction of said compound with a suitable acid or base using a conventional procedure.

When an optically active form of a compound of the formula I is required, it may be obtained, for example, by carrying out one of the aforesaid procedures using an optically

active starting material or by resolution of a racemic form of said compound using a conventional procedure, for example by the formation of diastereomeric salts, use of chromatographic techniques, conversion using chirally specific enzymatic processes, or by addition of temporary extra chiral group to aid separation.

5 As stated previously, the compounds of the formula I are inhibitors of the enzyme Factor Xa. The effects of this inhibition may be demonstrated using one or more of the standard procedures set out hereinafter:-

a) Measurement of Factor Xa Inhibition

10 An in vitro assay system is carried out based on the method of Kettner et al., J. Biol. Chem., 1990, 265, 18289-18297, whereby various concentrations of a test compound are dissolved in a pH7.5 buffer containing 0.5% of a polyethylene glycol (PEG 6000) and incubated at 37°C with human Factor Xa (0.001 Units/ml, 0.3 ml) for 15 minutes. The chromogenic substrate S-2765 (KabiVitrum AB, 20 µM) is added and the mixture is incubated at 37°C for  
15 20 minutes whilst the absorbance at 405 nm is measured. The maximum reaction velocity (V<sub>max</sub>) is determined and compared with that of a control sample containing no test compound. Inhibitor potency is expressed as an IC<sub>50</sub> value.

b) Measurement of Thrombin Inhibition

The procedure of method a) is repeated except that human thrombin (0.005 Units/ml) and the  
20 chromogenic substrate S-2238 (KabiVitrum AB, 7 µM) are employed.

c) Measurement of Anticoagulant Activity

An in vitro assay whereby human, rat or rabbit venous blood is collected and added directly to a sodium citrate solution (3.2 g/100 ml, 9 parts blood to 1 part citrate solution). Blood plasma is prepared by centrifugation (1000 g, 15 minutes) and stored at 2-4°C. Conventional  
25 prothrombin time (PT) tests are carried out in the presence of various concentrations of a test compound and the concentration of test compound required to double the clotting time, hereinafter referred to as CT<sub>2</sub>, is determined. In the PT test, the test compound and blood plasma are incubated at 37°C for 10 minutes. Tissue thromboplastin with calcium (Sigma Limited, Poole, England) is added and fibrin formation and the time required for a clot to  
30 form are determined.

d) An ex vivo Assay of Anticoagulant Activity

The test compound is administered intravenously or orally to a group of Alderley Park Wistar rats. At various times thereafter animals are anaesthetised, blood is collected and PT coagulation assays analogous to those described hereinbefore are conducted.

5 e) An in vivo Measurement of Antithrombotic Activity

Thrombus formation is induced using an analogous method to that described by Vogel et al., Thromb. Research, 1989, 54, 399-410. A group of Alderley Park Wistar rats is anaesthetised and surgery is performed to expose the vena cava. Collateral veins are ligated and two loose sutures are located, 0.7 cm apart, round the inferior vena cava. Test  
10 compound is administered intravenously or orally. At an appropriate time thereafter tissue thromboplastin (30 µl/kg) is administered via the jugular vein and, after 10 seconds, the two sutures are tightened to induce stasis within the ligated portion of vena cava. After 10 minutes the ligated tissue is excised and the thrombus therein is isolated, blotted and weighed.

15 (f) Rat Disseminated Intravascular Coagulation in vivo activity test

Fasted male Alderley Park rats (300-450 g) are pre-dosed by oral gavage (5 mls/kg) with compound or vehicle (5% DMSO/PEG200) at various times before being anaesthetised with Intraval® (120 mg/kg i.p.). The left jugular vein and the right carotid artery are exposed and  
20 cannulated. A 1 mL blood sample is taken from the carotid canular into 3.2% trisodium citrate. 0.5 mL of the whole blood is then treated with EDTA and used for platelet count determination whilst the remainder is centrifuged (5 mins, 20000g) and the resultant plasma frozen for subsequent drug level, fibrinogen or thrombin antithrombin (TAT) complex determinations. Recombinant human tissue factor (Dade Innovin Cat.B4212-50),  
25 reconstituted to the manufacturers specification, is infused (2 mL/kg/hr) into the venous canular for 60 minutes. Immediately after the infusion is stopped a 2 mL blood sample is taken and platelet count, drug level, plasma fibrinogen concentration and TAT complex are determined as before. Platelet counting is performed using at Coulter T540 blood analyser. Plasma fibrinogen and TAT levels are dertermining using a clotting assay (Sigma Cat.880-B)  
30 and TAT ELISA (Behring) respectively. The plasma concentration of the compound is bioassayed using human Factor Xa and a chromogenic substrate S2765 (Kabi), extrapolated from a standard curve (Fragmin) and expressed in Anti-Factor Xa units. The data is analysed

as follows; tissue factor-induced reductions in platelet count are normalised with respect to pre-dose platelet count and drug activity expressed as a percent inhibition of tissue factor-induced thrombocytopenia when compared to vehicle treated animals. Compounds are active if there is statistically significant ( $p < 0.05$ ) inhibition of TF-induced thrombocytopenia.

5 Example had an  $IC_{50}$  (Factor Xa) of  $0.007\mu M$  as measured in test a)

According to a further feature of the invention there is provided a pharmaceutical composition which comprises a heterocyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable diluent or carrier.

10 The composition may be in a form suitable for oral use, for example a tablet, capsule, aqueous or oily solution, suspension or emulsion; for topical use, for example a cream, ointment, gel or aqueous or oily solution or suspension; for nasal use, for example a snuff, nasal spray or nasal drops; for vaginal or rectal use, for example a suppository; for administration by inhalation, for example as a finely divided powder such as a dry powder, a  
15 microcrystalline form or a liquid aerosol; for sub-lingual or buccal use, for example a tablet or capsule; or for parenteral use (including intravenous, subcutaneous, intramuscular, intravascular or infusion), for example a sterile aqueous or oily solution or suspension. In general the above compositions may be prepared in a conventional manner using conventional excipients.

20 The amount of active ingredient (that is a heterocyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof) that is combined with one or more excipients to produce a single dosage form will necessarily vary depending upon the host treated and the particular route of administration. For example, a formulation intended for oral administration to humans will generally contain, for example, from 0.5 mg to 2 g of active  
25 agent compounded with an appropriate and convenient amount of excipients which may vary from about 5 to about 98 percent by weight of the total composition. Dosage unit forms will generally contain about 1 mg to about 500 mg of an active ingredient.

According to a further feature of the invention there is provided use of a heterocyclic derivative of the formula I, or a pharmaceutically-acceptable salt thereof, in the  
30 manufacture of a medicament for use in a method of treatment of the human or animal body by therapy.

The invention also includes the use of such an active ingredient in the production of a medicament for use in:-

- (i) producing a Factor Xa inhibitory effect;
- (ii) producing an anticoagulant effect;
- 5 (iii) producing an antithrombotic effect;
- (iv) treating a Factor Xa mediated disease or medical condition;
- (v) treating a thrombosis mediated disease or medical condition;
- (vi) treating coagulation disorders; and/or
- (vii) treating thrombosis or embolism involving Factor Xa mediated coagulation.

10 The invention also includes a method of producing an effect as defined hereinbefore or treating a disease or disorder as defined hereinbefore which comprises administering to a warm-blooded animal requiring such treatment an effective amount of an active ingredient as defined hereinbefore.

The size of the dose for therapeutic or prophylactic purposes of a compound of the  
15 formula I will naturally vary according to the nature and severity of the medical condition, the age and sex of the animal or patient being treated and the route of administration, according to well known principles of medicine. As mentioned above, compounds of the formula I are useful in the treatment or prevention of a variety of medical disorders where anticoagulant therapy is indicated. In using a compound of the formula I for such a purpose,  
20 it will generally be administered so that a daily dose in the range, for example, 0.5 to 500 mg/kg body weight is received, given if required in divided doses. In general lower doses will be administered when a parenteral route is employed, for example a dose for intravenous administration in the range, for example, 0.5 to 50 mg/kg body weight will generally be used. For preferred and especially preferred compounds of the invention, in general, lower  
25 doses will be employed, for example a daily dose in the range, for example, 0.5 to 10 mg/kg body weight.

Although the compounds of the formula I are primarily of value as therapeutic or prophylactic agents for use in warm-blooded animals including man, they are also useful whenever it is required to produce an anticoagulant effect, for example during the ex-vivo  
30 storage of whole blood or in the development of biological tests for compounds having anticoagulant properties.

The compounds of the invention may be administered as a sole therapy or they may be administered in conjunction with other pharmacologically active agents such as a thrombolytic agent, for example tissue plasminogen activator or derivatives thereof or streptokinase. The compounds of the invention may also be administered with, for example, a known platelet aggregation inhibitor (for example aspirin, a thromboxane antagonist or a thromboxane synthase inhibitor), a known hypolipidaemic agent or a known anti-hypertensive agent.

The invention will now be illustrated in the following Examples in which, unless otherwise stated:-

- 10 (i) evaporations were carried out by rotary evaporation in vacuo and work-up procedures were carried out after removal of residual solids by filtration;
- (ii) operations were carried out at room temperature, that is in the range 18-25°C and under an atmosphere of an inert gas such as argon;
- (iii) the end-products of the formula I have satisfactory microanalyses and their  
15 structures were confirmed by nuclear magnetic resonance (NMR) and mass spectral techniques. Chemical shift values were measured on the delta scale; the following abbreviations have been used: s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet;
- (iv) intermediates were not generally fully characterised and purity was assessed by thin layer chromatographic, infra-red (IR) or NMR analysis; and
- 20 (v) melting points were determined using a Mettler SP62 automatic melting point apparatus or an oil-bath apparatus; melting points for the end-products of the formula I were generally determined after crystallisation from a conventional organic solvent such as ethanol, methanol, acetone, ether or hexane, alone or in admixture.

## 25 Example 1

### 1-(5-Chloroindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine

A stirred suspension of 6-(4-pyridyl)nicotinic acid (400 mg, 2 mmol) in dimethylformamide, DMF, (10 ml) was treated with 1-(5-chloroindol-2-ylsulphonyl) piperazine (600 mg, 2 mmol, 30 1 mol eq.) and 1-(3-dimethylaminopropyl)-3-ethylcarbodi-imide hydrochloride (EDAC, 460 mg, 2.4 mmol, 1.2 mol eq.). After stirring overnight the solvent was removed *in vacuo* and the

- 19 -

residue chromatographed (Isolute 20g silica cartridge, eluting with dichloromethane containing 2.5% - 5% v/v of methanol) to yield 1-(5-chloroindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine as a colourless foam (680 mg). This was dissolved in dichloromethane/methanol mixture (40ml of 1:1) and treated with a saturated solution of HCl in methanol until acid to indicator paper (slight excess). The resulting solution of hydrochloride salt was evaporated to dryness and the residue boiled in 2-propanol (100ml, incomplete solution). Filtration and cooling gave 1-(5-chloroindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine hydrochloride as a colourless solid, (220 mg), <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 3.0-3.3 (broad d, 4H), 3.6-4.0 (broad d, 4H), 7.05 (s, 1H), 7.35 (dd, 1H), 7.5 (d, 1H), 7.8 (d, 1H), 8.1 (dd, 1H), 8.35 (d, 1H), 8.5 (m, 2H), 8.8 (d, 1H), 8.95 (d, 2H), 12.4 (s, 1H), signals were also present due to 2-propanol (0.5 mol equiv.); MS (M+H)<sup>+</sup> 481/483; mp 186 - 190 °C (not sharp).

The requisite 6-(4-pyridyl)nicotinic acid starting material was prepared as follows:

A solution of 1-[6-(4-pyridyl)3-pyridyl]-4-(tert.-butoxycarbonyl)-piperazine (3.7g, 10 mmol) and potassium carbonate (6.9g, 50 mmol) in methanol/water (90ml of a 2:1 mixture) (30 ml) was heated at reflux for 7 hrs. It was then cooled and neutralised with dilute HCl (50 ml of 2M), and some of the solvent removed in vacuo. More water was added and the resultant slurry left to stand for 2hrs. Filtration, washing with water and drying gave the above starting material (870 mg) which was used without further purification, <sup>1</sup>H NMR (d<sub>6</sub>-DMSO), 8.1 (d, 2H), 8.25 (d, 1H), 8.45 (dd, 1H), 8.75 (d, 2H), 9.2 (d, 1H), MS (M+H)<sup>+</sup> 201, (M-H)<sup>-</sup> 199.

1-[6-(4-Pyridyl)3-pyridyl]-4-(tert.-butoxycarbonyl)-piperazine was prepared as shown in Example 1 of PCT/GB98/02210.

1-(5-Chloroindol-2-ylsulphonyl) piperazine was prepared as shown in Example 3 of GB9809351.1.


**Example 2****1-(5-Bromoindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine**

By an exactly analogous method to that in Example 1, starting from 6-(4-pyridyl)nicotinic  
5 acid (400 mg, 2 mmol) and 1-(5-bromoindol-2-ylsulphonyl) piperazine (700 mg, 2 mmol, 1  
mol eq.), was prepared 1-(5-bromoindol-2-ylsulphonyl)-4-[6-(4-pyridyl)nicotinoyl] piperazine  
free base, as a colourless solid, (540 mg), <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 3.0-3.3 (broad d, 4H), 3.4-3.9  
(broad d, 4H), 7.0 (s, 1H), 7.45 (s, 2H), 7.95 (s, 1H), 8.0 (d, 1H), 8.1 (dd, 2H), 8.15 (d, 1H),  
8.75 (m, 3H), 12.4 (s, 1H), signals were also present due to DMF (1 mol equiv.); MS (M+H)<sup>+</sup>  
10 526/528.

The requisite 1-(5-bromoindol-2-ylsulphonyl) piperazine starting material was prepared in a  
manner analogous to that for the corresponding 5-chloro compound.

**Claims**

1. A compound of the formula I



wherein:

- A is an optionally substituted 5- or 6-membered monocyclic aromatic ring containing 1, 2 or 3 ring heteroatoms selected from oxygen, nitrogen and sulphur atoms;
- B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> are independently CH or a nitrogen atom, wherein the ring formed from B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> may optionally be substituted; with the proviso that at least one of B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is nitrogen;
- T<sub>1</sub> is CH or N;
- T<sub>2</sub> is CH or N; with the proviso that at least one of T<sub>1</sub> and T<sub>2</sub> is N;
- X<sub>1</sub> is SO, SO<sub>2</sub>, C(R<sub>4</sub>)<sub>2</sub> or CO when T<sub>1</sub> is CH or N; or in addition X<sub>1</sub> is O or S when T<sub>1</sub> is CH;
- and wherein each R<sub>4</sub> is independently hydrogen or (1-4C)alkyl;
- L<sub>1</sub> is (1-4C)alkylene or (1-3C)alkylenecarbonyl;
- R<sub>2</sub> is hydrogen or (1-4C)alkyl;
- R<sub>3</sub> is hydrogen or (1-4C)alkyl;
- or R<sub>2</sub> and R<sub>3</sub> are joined to form a (1-4C)alkylene or -CH<sub>2</sub>CO- group; wherein the ring formed by T<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, T<sub>2</sub> and L<sub>1</sub> is optionally substituted;
- X<sub>2</sub> is S(O)<sub>y</sub>, wherein y is one or two, C(R<sup>5</sup>)<sub>2</sub> or CO; and each R<sup>5</sup> is independently hydrogen or (1-4C)alkyl;
- Y is selected from hydrogen, halo, trifluoromethyl, trifluoromethoxy, cyano, hydroxy, amino, nitro, carboxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-(1-4C)alkylamino, (1-4C)alkoxycarbonyl, N-(1-4C)alkylcarbamoyl, N,N-di-(1-4C)alkylcarbamoyl, (2-4C)alkanoyl, (2-4C)alkanoylamino, hydroxy-(1-4C)alkyl, (1-4C)alkoxy-(1-4C)alkyl, carboxy-(1-4C)alkyl, (1-4C)alkoxycarbonyl-(1-4C)alkyl, carbamoyl-(1-4C)alkyl, N-(1-4C)alkylcarbamoyl-

- 22 -

(1-4C)alkyl and N,N-di-(1-4C)alkylcarbonyl-(1-4C)alkyl;

n is 1 or 2; and

B<sub>5</sub> and B<sub>6</sub> is selected from N or CH; with the proviso that at least one of B<sub>5</sub> and B<sub>6</sub> is N; and pharmaceutically acceptable salts thereof.

5

2. A compound of the formula I as claimed in claim 1 wherein A is a pyridyl, pyrimidinyl or pyridazinyl ring.

3. A compound of the formula I as claimed in claim 2 wherein A is 4-pyridyl, 2-pyridyl, 4-pyridazinyl, 3-pyrimidinyl, 4-pyrimidinyl or 3-pyridyl.

4. A compound of the formula I as claimed in claim 1 or 3 wherein A is unsubstituted or is substituted by one, two or three atoms or groups selected from halo, trifluoromethyl, cyano, amino, oxo, hydroxy, nitro, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)alkylamino or di-(1-4C)alkylamino.

5. A compound of the formula I as claimed in claim 1, 3 or 4 wherein the ring formed by B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is a pyridinediyl, wherein B<sub>1</sub>, or B<sub>3</sub> is a nitrogen atom, pyrimidinediyl, wherein B<sub>1</sub> and B<sub>2</sub> or B<sub>3</sub> and B<sub>4</sub> are nitrogen atoms or pyridazinediyl, wherein B<sub>1</sub>, B<sub>3</sub> and B<sub>4</sub> or B<sub>1</sub>, B<sub>2</sub> and B<sub>3</sub> are nitrogen atoms.

6. A compound of the formula I as claimed in claim 1, 3, 4 or 5 wherein the ring containing B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is unsubstituted or the ring containing B<sub>1</sub>, B<sub>2</sub>, B<sub>3</sub> and B<sub>4</sub> is substituted by one or two substituents selected from hydroxy, carboxy, (1-4C)alkoxycarbonyl or one of the following;

-(CH<sub>2</sub>)<sub>n</sub>-R, -(CH<sub>2</sub>)<sub>n</sub>-NRR<sub>1</sub>, -CO-R, -CO-NRR<sub>1</sub>, -(CH<sub>2</sub>)<sub>n</sub>-CO-R and -(CH<sub>2</sub>)<sub>n</sub>-CO-NRR<sub>1</sub>;

wherein n is 1 or 2;

30

R and R<sub>1</sub> are independently selected from hydrogen, (1-4C)alkyl, (2-4C)alkenyl,

(2-4C)alkynyl, hydroxy(1-4C)alkyl, carboxy(1-4C)alkyl and (1-4C)alkoxycarbonyl-(1-4C)alkyl or where possible R and R<sub>1</sub> may together form a 5- or 6-membered optionally substituted heterocyclic ring which may include in addition to the nitrogen atom to which R and R<sub>1</sub> are attached 1 or 2 additional heteroatoms selected from nitrogen, oxygen and sulphur.

5

7. A compound of the formula I as claimed in claim 1, 3, 4, 5 or 6 wherein the heterocyclic ring formed by R and R<sub>1</sub> is preferably selected from pyrrolidin-1-yl, imidazolin-1-yl, piperidin-1-yl, piperazin-1-yl, 4-morpholino and 4-thiomorpholino.

10 8. A compound of the formula I as claimed in claim 1, 3, 4, 5, 6 or 7 wherein the heterocyclic ring formed by R and R<sub>1</sub> is unsubstituted or the ring formed by R and R<sub>1</sub> is substituted by 1 or 2 substituents selected from oxo, hydroxy and carboxy.

9. A compound of the formula I as claimed in claim 1, 3, 4, 5, 6, 7 or 8 wherein when  
15 T<sub>1</sub> is CH or N then X<sub>1</sub> is CO, SO<sub>2</sub>, or CH<sub>2</sub> or when T<sub>1</sub> is CH then X<sub>1</sub> in addition is O or S.

10. A compound of the formula I as claimed in claim 1, 3, 4, 5, 6, 7, 8 or 9 wherein R<sub>2</sub> and R<sub>3</sub> are joined to form a C<sub>1-4</sub>alkylene group to form, together with T<sub>1</sub>, T<sub>2</sub> and L<sub>1</sub>, a heterocyclic ring selected from piperazine, piperidine and pyrrolidine.

20

11. A compound of the formula I as claimed in claim 1 or 3 wherein the heterocyclic ring formed by T<sub>1</sub>, T<sub>2</sub>, L<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> is unsubstituted or is substituted by one or two substituents selected from hydroxy, oxo, carboxy, (1-4C)alkoxycarbonyl or one of the following;

25

$-(CH_2)_n-R$ ,  $-(CH_2)_n-NRR_1$ ,  $-CO-R$ ,  $-CO-NRR_1$ ,  $-(CH_2)_n-CO-R$  and  $-(CH_2)_n-CO-NRR_1$ ;

wherein n is 1 or 2;

R and R<sub>1</sub> are independently selected from hydrogen, (1-4C)alkyl, (2-4C)alkenyl,  
30 (2-4C)alkynyl, hydroxy(1-4C)alkyl, carboxy(1-4C)alkyl and (1-4C)alkoxycarbonyl-

- 24 -

(1-4C)alkyl or where possible R and R<sub>1</sub> may together form a 5- or 6-membered optionally substituted heterocyclic ring which may include in addition to the nitrogen atom to which R and R<sub>1</sub> are attached 1 or 2 additional heteroatoms selected from nitrogen, oxygen and sulphur.

12. A compound of the formula I as claimed in claim 1, 3, 4, 5, 6, 7, 8, 9, 10 or 11

5 wherein X<sub>2</sub> is SO<sub>2</sub>, CH<sub>2</sub> or CO.

13. A compound of the formula I as claimed in claim 1, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 wherein Y is selected from hydrogen, halo, trifluoromethyl, trifluoromethoxy, cyano, hydroxy, amino, nitro, carboxy, carbamoyl, (1-4C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, (1-4C)alkoxy, 10 (2-4C)alkenyloxy, (2-4C)alkynyloxy, (1-4C)alkylthio, (1-4C)alkylsulphinyl, (1-4C)alkylsulphonyl, (1-4C)alkylamino, di-(1-4C)alkylamino and (1-4C)alkoxycarbonyl.

14. A compound of the formula I as claimed in any one of claims 1 to 12 for use in medical therapy.

15

15. A pharmaceutical composition which comprises a compound of the formula I, as defined in any one of claims 1 to 12 or a pharmaceutically-acceptable salt thereof, in association with a pharmaceutically-acceptable diluent or carrier.

20 16. Use of a compound of the formula I, or a pharmaceutically-acceptable salt thereof, as defined in any one of claims 1 to 12, in the manufacture of a medicament for use in a producing a Factor Xa inhibitory effect in a human.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00354

**A. CLASSIFICATION OF SUBJECT MATTER**  
 IPC 7 C07D401/14 A61K31/395

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
 IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 21188 A (ZENECA LTD.) 22 May 1998 (1998-05-22) claims	1, 15, 16
P, Y	WO 99 06371 A (ZENECA LTD.) 11 February 1999 (1999-02-11) * complete document *	1, 15, 16
P, Y	WO 99 57113 A (ZENECA LTD.) 11 November 1999 (1999-11-11) * complete document *	1, 15, 16

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*A\* document member of the same patent family

Date of the actual completion of the international search

11 May 2000

Date of mailing of the international search report

26/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/00354

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9821188	A	22-05-1998	AU 4874897 A	03-06-1998
			BR 9712672 A	26-10-1999
			CN 1235597 A	17-11-1999
			CZ 9901634 A	11-08-1999
			EP 0937048 A	25-08-1999
			NO 992230 A	07-05-1999
			PL 333241 A	22-11-1999
WO 9906371	A	11-02-1999	AU 8455798 A	22-02-1999
			NO 20000448 A	16-03-2000
WO 9957113	A	11-11-1999	AU 3620699 A	23-11-1999